

## Biological Control of Potato Bacterial Wilt Diseases

Ali Ahmed Waheed<sup>1,2</sup>, Mohamed Ebrahim<sup>1</sup>, Afaf, Z.A. El-Meneisy<sup>1</sup>, Waffa, M. Abd El-Sayid<sup>1</sup>

<sup>1</sup>Department of Plant Pathology, Faculty of Agriculture, Ain Shams University, Cairo, Egypt

<sup>2</sup>Faculty of Agriculture, University of Al-Qadisiyah, Iraq.

Email: [Aliwaheed\\_2020@agr.asu.edu.eg](mailto:Aliwaheed_2020@agr.asu.edu.eg)<sup>1,2</sup>, [Mohamed\\_gaber@agr.asu.edu.eg](mailto:Mohamed_gaber@agr.asu.edu.eg),  
[afaf\\_abdelaziz@agr.asu.edu.eg](mailto:afaf_abdelaziz@agr.asu.edu.eg), [wafaa@hotmail.com](mailto:wafaa@hotmail.com)

\*Corresponding author's E-mail: [Aliwaheed\\_2020@agr.asu.edu.eg](mailto:Aliwaheed_2020@agr.asu.edu.eg)

| Article History                                                          | Abstract                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           |
|--------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Received: 06 June 2023<br>Revised: 05 Sept 2023<br>Accepted: 14 Oct 2023 | <i>Potato (Solanum tuberosum L.) is considered one of the four major and important food crops around the world. This study planned to control bacterial wilt disease of potato using some bio-agents which isolated and identified from potato plant soil. Twenty isolates were selected and identified as the following three isolates belonging to Streptomyces spp. (Streptomyces antibioticus (SA1), S. albus (SA2), and S. mutabilis (SM1); five isolates belong to pseudomonas species (Pseudomonas fluorescence (PsF1), P. aeruginosa (PsA1), P. putida (PsP1), P. alcaligenes (PsA2) and P. pseudoalcaligones (PsP2)) and twelve isolates were belonging to Bacillus spp. (Bacillus subtilis (BS1-8), B. cereus(BC1), B. badius (BB1-2), B. pumilus (BP1). In vitro, these isolates were examined against the growth R. solanacearum bacterium, where some isolates (BS3, PsF1, BS8, BS6, SM1, BS5, and BS4) were the most effective compared with other isolates. Bacillus subtilis (BS8); Pseudomonas fluorescence (PsF1) and Streptomyces mutabilis (SM1) isolates were selected as bio-agents to control potato bacterial wilt disease under in vivo condition, where these isolates led to reduced disease severity and to increase potato yield compared with the control. The application of bio-agents as drench treatment was more effective than tuber treatment, and isolates of S. mutabilis (SM1) and B. subtilis (BS8) were more effective than Pseudomonas fluorescence (PsF1) isolate.</i> |
| CC License<br>CC-BY-NC-SA 4.0                                            | <b>Keywords:</b> Potato, Solanum tuberosum L., Bacterial wilt disease, Bio-agents, Ralstonia solanacearum, Biological control                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |

### 1. Introduction

*Ralstonia solanacearum*, the causative of bacterial wilt (brown rot of potato) disease is widely spread in tropical and subtropical areas. This bacterium is a unique plant pathogen according to its economic and scientific value Mansfield et al., (2012). Brown rot is the second most important tropical and subtropical potato disease after late blight. Champoiseau et al., (2009). Brown rot disease, the most important potato bacterial disease in Egypt, was announced many years ago. Khairy et al., (2021). It is challenging to discover an effective method of controlling potato bacterial wilt because of the pathogen's persistence and ability to spread using irrigation water, soil, surface water, agricultural machinery, and infected organic material Janse, (1996) and Tan et al., (2016).

Biological control is supposed to preserve environmental quality by minimizing dependency on chemical inputs and maintaining sustainable management practices Bara and Jeffries, (1995). Soesanto et al., (2013) reported the successful antagonistic *Bacillus* sp. and *Pseudomonas fluorescens* isolated from the rhizosphere to control bacterial wilt on potatoes by delaying the incubation period suppressing disease intensity decreasing the final pathogenic population and inducing plant resistance. Plant growth-promoting Rhizobacteria (PGPR) are recognized for growth promotion and disease

control Maji and Chakrabarty, (2014) and Apastambh et al., (2016). Many chemical, biological, and physical tools were used to control *R. solanacearum* bacterium. Biological control agents are hopeful eco-friendly tools alternative to pesticides to manage diseases that are beneficial in protecting crop plants and safeguarding food as well He et al., (2021). Suppression mechanisms are typically attributed to the antibacterial metabolites produced by biological control agents or those present in natural products and increasing host resistance to the pathogen. Nion and Toyota, (2015).

Cultural and biological control of bacterial diseases were alternative solutions with some promising success Michel and Mew, (1998) Ran et al., (2005). Bacterial wilt disease cannot be controlled by chemical means. Up till now, no effective chemical products are available for controlling “bacterial wilt” (*R. solanacearum*). Among the several methods of disease management, biocontrol plays an important role in disease control and may help the development of alternative management measures being integrated with other practices for effectiveness. Control and for minimizing environmental Pollution due to the use of chemical pesticides Lwin and Ranamukhaarachchi, (2006) and Achari and Ramesh, (2014). The present work was planned to isolate and identify some antagonistic microorganism and their effects on the growth of *R. solanacearum*, the severity of Potato bacterial wilt, and potato yield under artificial inoculation conditions.

## **2. Materials and Methods**

### **Isolation and identification of potato bacterial wilt pathogen**

Samples of potato tubers with typical disease symptoms of bacterial wilt were collected from different districts in some governorates. Isolation procedures were carried out using tetrazolium chloride (TZC) medium and South Africa (SMSA) described by Engelbrecht (1994) and Elphinstone (1996). Collected isolates were identified using cultural, and morphological characteristics, and biochemical and physiological tests were studied using the methods described by Schaad et al., (2001); and McCarter (1976). Immunofluorescence antibody staining (IFAS) test was applied to confirm their identification in Potato Brown Rot Project, Dokki, Giza, Egypt (2022). Positive isolates were inoculated in tomato seedlings to verify their pathogenic potential (Janse, 1988 and Wenneker et al., 1999).

### **Isolation of antagonistic microorganisms from potato rhizosphere**

Ten grams of potato plants rhizosphere soil were collected from infested fields with bacterial wilt disease. Each sample was placed in flask (250 ml) containing 90 ml of sterile distilled water and shaken for 30 min. using horizontal shake. Serial dilution was prepared and one ml of  $10^4$ ,  $10^5$  and  $10^7$  dilution were poured on nutrient agar medium for *Bacillus* spp according to (Lelliott and Stead, 1987), on king's B medium for *Pseudomonas* fluorescents species (King et al., 1954) and on starch nitrate agar medium for actinomycetes species (Shirling and Gottlieb, 1966). These plates were gently rotated and incubated at  $27 \pm 1^\circ\text{C}$  for 3, 7 and 14 days, respectively. Selected colonies were transferred from mixed culture of the plates onto respective agar plates for purification by streak plate technique and the plates were incubated at  $27 \pm 1^\circ\text{C}$  for 2-7 days. After purification, plates containing pure cultures were stored at  $4^\circ\text{C}$  until further studies. Four plates were used as replicates for each treatment.

### **Identification of antagonistic bacteria and actinomycetes isolates**

Identification of the antagonistic bacteria (*Bacillus* spp and *Pseudomonas* species) was made through morphological as well as biochemical characteristics according to the method described by Fahy and Persley (1983). Actinomycetes isolates were identified as described by Cross (1989); Locci (1989) and Williams, et al. (1989). All identification procedures were carried out in the regional center of Mycology and Biotechnology, Al-Azhar University, Cairo, Egypt.

The antagonistic effect of isolated bacteria and actinomycetes against *R. solanacearum* bacterium, in vitro: The antagonistic effect of bacteria and actinomycetes against the growth of *R. solanacearum* bacterium, using the technique described by Madigan, et al. (1997). The plates were prepared and inoculated with the tested bacteria or actinomycetes cultures by a single streak of inoculum at a certain distance (3 cm) from the center of the petri dish and incubated at  $27 \pm 1^\circ\text{C}$  for 4 days for actinomycetes and 2 days for the tested bacteria. Later, the plates were seeded with *R. solanacearum* by a single streak starting from the center of the plate and directed to the tested organism (bacteria or

Actinomycetes) at a 90° angle to the tested organism. Antagonism was measured by determining the length of the inhibition zones (mm), and different inhibition zones were recorded. The antagonistic effect of isolated bacteria and actinomycetes against *R. solanacearum* bacterium, *in vivo*:

The virulent isolate of *Ralstonia solanacearum* (RS3) isolated from an infected potato tuber was grown on sucrose peptone agar medium for 48 h at 28°C. Bacterial growths were suspended in sterilized tap water and the population density of the bacterium was adjusted to give  $6 \times 10^8$  colony-forming units (CFU/ml) using a spectrophotometer (OD600 = 0.3). The sterilized clay soil was distributed in sterilized clay pots 30 cm in diameter (7 Kg soil/pot) and separately infested with the bacterial inoculum to give a final concentration of  $10^7$  CFU/g dry soil. The infestation procedure was made five days before planting according to Michel and Mew (1998). Antagonistic isolates were grown on an NA medium for 48 h for (bacteria) and 7 days for (actinomycetes). A concentration of  $10^9$  CFU/ml was prepared for each antagonistic bacterium and actinomycetes using a Spectrophotometer (OD600= 1) according to Ciampi and Sequeira (1980). Antagonists were applied singly as soil or tuber treatment. In soil treatment, each pot was treated with 500 ml of the antagonistic isolate at the time of sowing. Meanwhile, in tuber treatment, potato tubers (cv, Nicola) were soaked in the suspensions of the antagonistic bio-agents plus 0.1% methylcellulose (1:1 v/v) for 60 minutes before sowing in infested soil. Four pots were used as replicates for each treatment. Twelve replicates with only the pathogen were used as control treatments. All pots were kept under outdoor conditions.

### Disease assessment

Disease severity was assessed after 70 and 90 days after planting. The severity of bacterial wilt was calculated as a percentage of sprouts showing wilt symptoms per plant and the disease index (%) was calculated from disease rating for individual plants (Kempe and Sequeira 1983) using the scale based on the visual observation of the percentage of foliage wilt (0 = no visible symptoms, 1 = 1- 25% wilt, 2= 26- 50%, 3 = 51-75% wilt, 4= 76- 100% wilt and 5 = dead plants. The percentage of disease index (DI%) was calculated by the following formula:

$$DI = \frac{\sum RT}{5N} \times 100$$

**Where:** T = Total number of plants with each category.

**R** = Disease severity scale (R = 1, 2, 3, 4 and 5)

**N** = Total number of tested plants.

Disease reduction percentages (DRP) were calculated from disease index using the following formula:

$$PDR = \frac{DI_{ck} - DI_{tr}}{DI_{ck}} \times 100$$

**Where:**  $DI_{ck}$  = Disease index in check treatment.

$DI_{tr}$  = Disease index in treated treatment.


### Statistical analysis

The determination of statistical difference was based on a comparison of the disease severity and the yield. First, the initial conditions of validation were tested to determine which approach, parametric or nonparametric, should be used. If conditions of normal residues distribution and homoscedasticity were fulfilled, a parametric approach based on an ANOVA followed by pairwise multiple comparisons (ad hoc test of Duncan's method) was performed. Otherwise, a nonparametric approach based on a Kruskal-Wallis test was used. The analyses were performed, using the Excel Add-in XLSTAT software program (Addinsoft 2021).

### 3. Results and Discussion

#### Isolation and identification of the causal organism from infected potato tubers

Collected isolates were identified as *Ralstonia solanacearum* bacterium according to morphological, cultural, physiological, and biochemical characteristics. This identification was performed with the immunofluorescent staining system in Potato Brown Rot Project. (fig. 1).



**Fig. (1):** Cell morphology of *R. solanacearum* in the serological immunofluorescent antibody staining (IFAS) test where cells are short rods and stained evenly as bright green fluorescent under an IF microscope

#### Identification of antagonistic microorganisms from soil potato plants

Twenty isolates were selected and identified according to cultural, morphological, and biochemical properties (Table 1-3). Data in Table (1) revealed that three isolates belonged to *Streptomyces* spp, where these isolates were identified as the following *Streptomyces antibioticus* (SA1), *S. albus* (SA2), and *S. mutabilis* (SM1). Also, five isolates were belonging to *pseudomonas* species, where these isolates were identified as the following *Pseudomonas fluorescence* (PsF1), *P. aeruginosa* (PsA1), *P. putida* (PsP1), *P. alcaligenes* (PsA2) and *P. pseudoalcaligones* (PsP2) (Table 2). However, twelve isolates belonging to *Bacillus* spp as the following *Bacillus subtilis* (BS1-8), *B. cereus* (BC1), *B. badius* (BB1-2), *B. pumilus* (BP1) (Table 3).

**Table (1):** Characterizations of isolated antagonistic actinomycetes (*Streptomyces* spp) and their ability to carbon sources utilization

| Characteristic           | Code isolate of actinomycetes |                      |                      |
|--------------------------|-------------------------------|----------------------|----------------------|
|                          | SA1                           | SA2                  | SM1                  |
| Sporophore morphology    | Straight to flexible          | Straight to flexible | Straight to flexible |
| Spore surface            | smooth                        | spiny                | Smooth               |
| Growth                   | Well                          | Well                 | moderate             |
| Color of aerial mycelium | Gray                          | Grayish brown        | white                |
| Color of growth          | Brownish gray                 | brownish yellow      | Non-distinctive      |
| Soluble pigment          | Negative                      | Negative             | Negative             |
| Cellulose decomposition  | negative                      | Negative             | Negative             |
| Gram stain               | +                             | +                    | +                    |
| Sporulation              | +                             | +                    | +                    |
| Cell shape               | Hyphae                        | Hyphae               | Hyphae               |
| KOH test                 | -                             | -                    | -                    |
| Motility                 | -                             | -                    | -                    |
| d. Glucose               | +                             | +                    | +                    |
| D. Fructose              | +                             | -                    | +                    |
| D. xylose                | +                             | -                    | +                    |
| L. rhaminose             | +                             | +                    | +                    |

|                |                        |                 |                     |
|----------------|------------------------|-----------------|---------------------|
| Rafinose       | -                      | +               | -                   |
| Sucrose        | -                      | +               | +                   |
| Glycerol       | +                      | +               | +                   |
| Starch         | +                      | +               | +                   |
| Identification | <i>S. antibioticus</i> | <i>S. albus</i> | <i>S. mutabilis</i> |

**Table (2):** Morphological and biochemical Characteristics of antagonistic bacteria (*Pseudomonas* spp.)

| Characteristic       | Code isolate of bacteria |                      |                  |                       |                             |
|----------------------|--------------------------|----------------------|------------------|-----------------------|-----------------------------|
|                      | PsF1                     | PsA1                 | PsP1             | PsA2                  | PsP2                        |
| Cell shape           | Short rods               | Short rods           | Short rods       | Short rods            | Short rods                  |
| Gram staining        | -                        | -                    | -                | -                     | -                           |
| Sporulation          | -                        | -                    | -                | -                     | -                           |
| Arginine dihydrolase | +                        | +                    | +                | +                     | +                           |
| Levan production     | +                        | -                    | -                | +                     | +                           |
| Gelatin hydrolysis   | +                        | +                    | +                | +                     | +                           |
| Oxidase              | +                        | +                    | -                | +                     | +                           |
| Catalase             | +                        | +                    | -                | -                     | +                           |
| Starch hydrolase     | -                        | -                    | -                | -                     | -                           |
| Indole production    | -                        | -                    | -                | +                     | +                           |
| Pigment on KB medium | +                        | +                    | +                | -                     | -                           |
| Utilization of:      |                          |                      |                  |                       |                             |
| Glucose              | +                        | +                    | +                | -                     | -                           |
| Trehalose            | +                        | -                    | -                | -                     | -                           |
| D-Mannose            | +                        | -                    | +                | +                     | +                           |
| Sucrose              | +                        | -                    | -                | +                     | +                           |
| Identification       | <i>P. fluorescens</i>    | <i>P. aeruginosa</i> | <i>P. putida</i> | <i>P. alcaligenes</i> | <i>P. pseudoalcaligones</i> |

**Table 3:** Morphological and biochemical Characteristics of antagonistic bacteria (*Bacillus* spp.)


| Characteristic       | Code isolate of bacteria |           |           |           |           |           |           |           |           |           |           |           |
|----------------------|--------------------------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|-----------|
|                      | BS1                      | BS2       | BS3       | BS4       | BS5       | BS6       | BS7       | BS8       | BC1       | BB1       | BB2       | BP1       |
| Cell shape           | Long rods                | Long rods | Long rods | Long rods | Long rods | Long rods | Long rods | Long rods | Long rods | Long rods | Long rods | Long rods |
| Gram staining        | +                        | +         | +         | +         | +         | +         | +         | +         | +         | +         | +         | +         |
| Sporulation          | +                        | +         | +         | +         | +         | +         | +         | +         | +         | +         | +         | +         |
| Arginine dihydrolase | +                        | +         | +         | +         | +         | +         | +         | +         | +         | +         | +         | +         |
| Levan production     | -                        | -         | -         | -         | -         | -         | -         | -         | -         | -         | -         | -         |
| Gelatin hydrolysis   | +                        | +         | +         | +         | +         | +         | +         | +         | +         | +         | +         | -         |



|                   |                    |                    |                    |                    |                    |                    |                    |                    |                  |                  |                  |                   |
|-------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|--------------------|------------------|------------------|------------------|-------------------|
| Oxidase           | -                  | -                  | -                  | -                  | -                  | -                  | -                  | -                  | +                | +                | +                | +                 |
| Catalase          | +                  | +                  | +                  | +                  | +                  | +                  | +                  | +                  | +                | +                | +                | +                 |
| Starch hydrolase  | +                  | +                  | +                  | +                  | +                  | +                  | +                  | +                  | +                | -                | -                | -                 |
| Indole production | -                  | -                  | -                  | -                  | -                  | -                  | -                  | -                  | -                | -                | -                | -                 |
| Utilization of:   |                    |                    |                    |                    |                    |                    |                    |                    |                  |                  |                  |                   |
| Glucose           | +                  | +                  | +                  | +                  | +                  | +                  | +                  | +                  | +                | -                | -                | -                 |
| Arabinose         | +                  | +                  | +                  | +                  | +                  | +                  | +                  | +                  | -                | -                | -                | -                 |
| Identification    | <i>B. subtilis</i> | <i>B. subtilis</i> | <i>B. subtilis</i> | <i>B. subtilis</i> | <i>B. subtilis</i> | <i>B. subtilis</i> | <i>B. subtilis</i> | <i>B. subtilis</i> | <i>B. cereus</i> | <i>B. badius</i> | <i>B. badius</i> | <i>B. pumilus</i> |

### Effect of isolated microorganisms on the growth of *R. solanacearum*, in vitro

Twenty isolates of antagonistic microorganisms were examined against the growth *R. solanacearum* bacterium, in vitro (fig. 2). Five isolates of *Bacillus subtilis* (BS3, BS8, BS6, BS5, and BS4); one isolate of *Pseudomonas fluorescens* (PsF1) and one isolate of *Streptomyces mutabilis* (SM1) was the most effective to inhibit the growth *R. solanacearum* bacterium, where the diameter of inhibition zone was 10.9, 10.9, 10.7, 10.4, 10.3, 10.9 and 10.6 mm, respectively. Meantime, three isolates of *B. subtilis* (BS1, BS7, and BS2); one isolate of *B. cereus* (BC1); three isolates of *Pseudomonas* spp (*P. alcaligenes* (PsA2); *P. pseudoalcaligenes* (PsP2) and *P. aeruginosa* (PsA1)) and two isolates of *Streptomyces* spp, (*S. antibioticus* (SA1) and *S. albus* (SA2)) were moderately effective, where the inhibition zone was 5.9, 5.5, 5.3, 5.6, 5.4, 5.4, 5.1, 5.9, and 5.4 mm, respectively. Meanwhile, three isolates of *Bacillus* spp. (*B. pumilus* (BP1) and two isolates of *B. badius* (BB1, BB2)) and one isolate of *P. putida* (Psp1) were less effective, where the inhibition zone was 0.9, 0.5, 0.3, and 0.5 mm, respectively.




**fig. (2):** Effect of some bio-agents on the growth of potato bacterial wilt bacterium (*Ralstonia solanacearum*), in vitro


**Effect of some bio-agents on bacterial wilt disease of potato and the total yield**

*Bacillus subtilis* (BS8); *Pseudomonas fluorescens* (PsF1) and *Streptomyces mutabilis* (SM1) isolates were applied as bio-agents to control potato bacterial wilt disease under artificial inoculation conditions (fig. 3-7). Data in figures (3 and 4) showed that examined bio-agents decreased the severity of bacterium wilt disease in potato compared with the control treatment. Likewise, the application of bio-agents as drench treatment was more effective than tuber treatment against the disease, where efficiency treatment ranged from 9.2 to 16.4% and from 5.8 to 15.2%, respectively. While, isolates of *S. mutabilis* (SM1) and *B. subtilis* (BS8) were more effective to control the disease than *Pseudomonas fluorescens* (PsF1) isolate, where efficiency treatment was 5.4 - 16.4, 5.0 - 16.0 and 5.8 - 13.6 %, respectively.


However, results in figure (5-6) appeared that the application of *B. subtilis* (BS8); *S. mutabilis* (SM1), and *P. fluorescens* (PsF1) isolates as bio-agents led to reducing the severity of infected tubers and increased potato yield (number and weight tubers) compared with the control treatment. While, drench treatment was more effective than tubers treatment, where efficiency for controlling infected tuber was 10.3 - 27.5% and 3.3 - 22.4%, and potato tubers number was 2.5 - 3.3 for 2.3 -3.0 tuber/plant and 44.30 - 65.05 and 40.64 - 60.60 g/plant, respectively. Meanwhile, *B. subtilis* (BS8) and *S. mutabilis* (SM1) isolates were more effective than *Pseudomonas fluorescens* (PsF1) isolate, where efficiency to control infected tuber was 22.4 - 27.5; 16.3 - 21.8% and 3.3 -10.3 % and the yield were 3.0 - 3.3; 2.5 - 2.7 and 2.3 - 2.5 tuber/plant and 60.60 - 65.05; 49.53 - 52.56 and 40.64 - 44.30 g/plant, respectively.




**Fig 3:** The efficiency of bio-agents on the severity of potato bacterial wilt disease, using two methods of treatment, after 70 days from artificial inoculation.



**Fig 4:** The efficiency of bio-agents on the severity of potato bacterial wilt disease, using two methods of treatment, after 90 days from artificial inoculation.



**Fig 5:** The efficiency of bio-agents on the severity of infected tubers (%), using two treatment methods, after 105 days from planting, under artificial inoculation



**Fig 6:** Effect of some bio-agents on the yield (A. Number of tubers, B. The weight of tubers/plant), using two treatment methods, after 105 days from planting, under artificial inoculation.



#### 4. Discussion:

Twenty isolates of antagonistic microorganisms were isolated and identified from rhizosphere potato plants as the following: Three isolates belong to *Streptomyces* species (*Streptomyces antibioticus* (SA1), *S. albus* (SA2), and *S. mutabilis* (SM1)). Also, five isolates belong to *Pseudomonas* species (*Pseudomonas Fluorescent* (PsF1), *P. aeruginosa* (PsA1), *P. putida* (PsP1), *P. alcaligenes* (PsA2) and *P. pseudocaligones* (PsP2). Meanwhile, twelve isolates belong to *Bacillus* species (eight isolates of *B. subtilis* (BS1-8), two isolates of *Bacillus badius* (BB1-2), *B. cereus* (BC1) and *B. Pumilus* (BP1). Soesante *et al.*, (2013) mentioned successfully isolated antagonistically *Bacillus* spp. and *Pseudomonas fluorescens* from rhizosphere Potato plants with infection by bacterial wilt disease and delaying incubation Period. Zhu *et al.*, (2021) mentioned that *Bacillus* species have a long history in biocontrol and crop growth-promoting applications. *Pseudomonas aeruginosa* isolate showed higher antifungal activity and a lower minimum inhibitory concentration Shiakh *et al.*, (2014).

Some plant-associated bacteria are classified as beneficial microorganisms based on their effects on plant performance, and free-living plant growth-promoting rhizobacteria (PGPR) thrive freely in the rhizosphere soil (Labuschagne *et al.*, (2011), Shaikh *et al.*, (2018)). Antagonistic bacteria from the genera *Pseudomonas*, *Bacillus* and *Streptomyces*, account for the majority of rhizosphere microorganisms commonly used in biocontrol. Ciancio *et al.*, (2019).

Isolates of antagonistic microorganisms appeared differential effective against the growth of *R. solanacearum* bacterium *in vitro*. *Bacillus subtilis* (BS8), *P. fluorescent* (PsF1), and *S. mutabilis* (SM1) isolates were the most effective compared with other isolates. These isolates were selected as bio agents to control bacterial wilt disease of potato, under artificial inoculation conditions. Ali *et al.*, (2013) found that plant growth-promoting rhizobacteria (PGPR) excretion different antibacterial substances in the culture medium. These substances reduce the growth of the bacteria and formed an inhibition zone. Balabel *et al.*, (2013) applied some antagonistic species against five isolates of *Ralstonia solanacearum* bacterium which showed varying antagonistic potential. Abd El Rahman and Shaheen (2016) found that *Bacillus subtilis*, *B. cereus*, *Pseudomonas fluorescens*, and *Rhizobium phaseoli* affect the growth of *Ralstonia solanacearum*. *In vitro*, where *B. subtilis* was the most effective, and *P. fluorescens* had the most growth potential and great inhibition on King's B medium. Arfaoui *et al.*, (2006) and Aliye *et al.*, (2008) reported that antagonism of *Bacillus* Spp, *P. fluorescens* and *Rhizobium* spp against the different plant pathogens, *In vitro*.

Selected antagonistic microorganism isolates decreased the severity of potato bacterial wilt and increased potato yield compared with the control. Likewise, the application of bio agents as soil drench treatment was more effective than their application as tubers treatment. *Streptomyces mutabilis* (SM1) and *Pseudomonas fluorescens* (PsF1) isolates were highly effective on disease severity and the yield compared with *Bacillus subtilis* (BS8) isolate. Wagih (1991) and Sunaina *et al.*, (1997) revealed that potato bacterial wilt (brown rot) disease reduced significantly after bacterization of healthy tuber potatoes with *Bacillus cereus*, *B. subtilis*, and with a virulent strain of *R. solanacearum* under field conditions. Aspiras and Cruz (1985) mentioned that *P. fluorescens* was highly effective in reducing bacterial wilt in tomato and potato under experimental conditions by aggressively colonizing the roots of young plants and preventing entry of *R. solanacearum*. Buyer, and Leong, (1986) and Ran *et al.*, (2005) found that inhibition of the pathogen by *P. fluorescens* was attributed to the siderophore pseudobactin, depriving root colonizing microorganisms, including plant pathogens of  $\text{Fe}^{3+}$  and inducing systemic resistance.

Application of *Bacillus* sp., *B. subtilis*, and *P. fluorescens* effectively controlled brown rot pathogen and increased potato yield Atia *et al.*, (2010). Rosyidah *et al.*, (2013) reported that a single application of *P. fluorescens* or combined application of *Streptomyces* spp and *Trichoderma viride* plus *Streptomyces* spp was capable reducing disease incidence, disease intensity, and, the population of *R. solanacearum*. Bacterial isolates' ability to produce antagonistic and growth-promoting substances modified their ability to increase plant growth and inhibited other organisms. Compant *et al.*, (2005). Perez-Garcia *et al.*, (2011) mentioned that promising achievements in terms of biological control have emerged, especially after the successful use of certain antagonistic biocontrol agents (BCAs), in particular *Pseudomonas* spp *Bacillus* spp *Burkholderia* spp and *Trichoderma* spp against pathogens causing foliar and soil-borne diseases. Alizadeh *et al.*, (2013) showed that biopesticides are available

inside the rhizosphere to combat pests and microbial diseases due to the close connection of root-colonizing probiotic microorganisms with plant host cells.

## 5. Conclusion

This study planned to control the bacterial wilt disease of potato using some bio-agents isolated and identified from potato plant soil. The antagonistic effect of 20 isolates obtained in vitro was tested against isolates of *R. solanacearum*, and the results showed that five isolates of *B. subtilis* (BS4, BS5, BS6, BS8, BS3), *P. fluorescence* bacterium, a single isolate (PsF1), and one isolate belonging to *S. mutabilis* (SM1) significantly reduced the growth of *R. solanacearum* compared with other isolates. Among the tested isolates, these particular ones showed the highest effectiveness level compared to the other isolates. The study employed *B. subtilis* (BS8), *P. fluorescence* (PsF1), and *S. mutabilis* (SM1) isolates as bio-agents for the management of potato bacterial wilt disease under greenhouse conditions. The results demonstrated that the application of these isolates resulted in a decrease in disease severity and an increase in potato yield when compared to the control group. The application of bio-agents as soil treatments was more effective than tuber treatment in decreasing the disease severity and increasing the potato yield.

## References:

- Abd El-Rahman, A. F., & Shaheen, H. A. (2016). Biological control of the brown rot of potato, *Ralstonia solanacearum* and effect of bacterization with antagonists on promotion of potato growth. *Egyptian J Biol Pest Control*, 26(4), 733-739.
- Achari, G. A., & Ramesh, R. (2014). Diversity, biocontrol, and plant growth promoting abilities of xylem residing bacteria from solanaceous crops. *International journal of microbiology*, 2014.
- Addinsoft (2021) Statistical software for excel; XLSTAT statistical and data analysis solution, version 2021, 2, Addinsoft, Paris. <https://www.xlstat.com/en/>.
- Ali Shahid, A., Yasin, S., Inam-ul-Haq, M., Ali, M. and Saleem Haider, M. (2013). "Use of Rhizobacteria for the Management of Soft Rot Disease of Potato" Athens: ATINER'S Conference Paper Series, No: AGR2013-0770
- Aliye, N., Fininsa, C., & Hiskias, Y. (2008). Evaluation of rhizosphere bacterial antagonists for their potential to bioprotect potato (*Solanum tuberosum*) against bacterial wilt (*Ralstonia solanacearum*). *Biological Control*, 47(3), 282-288.
- Alizadeh, O., Azarpanah, A., & Ariana, L. (2013). Induction and modulation of resistance in crop plants against disease by bioagent fungi (arbuscular mycorrhiza) and hormonal elicitors and plant growth promoting bacteria. *International Journal of Farming and Allied Sciences*, 2, 982-998.
- Apastambh, A. R., Tanveer, K., & Baig, M. M. V. (2016). Isolation and characterization of plant growth promoting Rhizobacteria from banana rhizosphere. *Int. J. Curr. Microbiol. App. Sci*, 5(2), 59-65.
- Arfaoui, A., Sifi, B., Boudabous, A., Hadrami, I. E., & Cherif, M. (2006). Identification of Rhizobium isolates possessing antagonistic activity against *Fusarium oxysporum* f. sp. *ciceris*, the causal agent of *Fusarium* wilt of chickpea. *Journal of Plant Pathology*, 67-75.
- Aspiras, R. B., & Cruz, A. R. (1985). Potential biological control of bacterial wilt in tomato and potato with *Bacillus polymyxa* FU6 and *Pseudomonas fluorescens*. *Bacterial wilt disease in Asia and the South Pacific*, 61, 89-92.
- Atia, M.M., M.M.A. Tohamy, F.F. Gabriel and H. A. Mater (2010). Integrated management of potato brown rot. *Proc.of the Sixth Int. Conf. Susation Agric., Fayoum Univ.*, 275- 294.
- Balabel, N. M., Messiha, N. A., & Farag, N. S. (2013). New findings on biological control trials of potato brown rot with antagonistic strains of *Bacillus circulans*, in Egypt. *Plant Pathology Journal (Faisalabad)*, 12(1), 11-18.
- Barea, J. M., & Jeffries, P. (1995). Arbuscular mycorrhizas in sustainable soil-plant systems. *Mycorrhiza: structure, function, molecular biology and biotechnology*, 521-560.
- Buyer, J. S., & Leong, J. (1986). Iron transport-mediated antagonism between plant growth-promoting and plant-deleterious *Pseudomonas* strains. *Journal of Biological Chemistry*, 261(2), 791-794.
- Champoiseau, P. G., Jones, J. B., & Allen, C. (2009). *Ralstonia solanacearum* race 3, biovar 2 causes tropical losses and temperate anxieties. *Plant Health Progress*, 10(1), 35.
- Ciampi, L., & Sequeira, L. (1980). Multiplication of *Pseudomonas solanacearum* in resistant potato plants and the establishment of latent infections. *American Potato Journal*, 57, 319-329.
- Ciancio, A., Pieterse, C. M., & Mercado-Blanco, J. (2019). Harnessing useful rhizosphere microorganisms for pathogen and pest biocontrol. *Frontiers in microbiology*, 10, 1935.

- Compant, S., Duffy, B., Nowak, J., Clément, C., & Barka, E. A. (2005). Use of plant growth-promoting bacteria for biocontrol of plant diseases: principles, mechanisms of action, and future prospects. *Applied and environmental microbiology*, 71(9), 4951-4959.
- Cross, T. (1989). Growth and examination of actinomycetes-some guidelines. *Bergey's manual of systematic bacteriology*, 4, 2340-2343.
- Elphinstone, J. G. (1996). Survival and possibilities for extinction of *Pseudomonas solanacearum* (Smith) Smith in cool climates. *Potato Research*, 39, 403-410.
- Engelbrecht, M. C. (1994). Modification of a semi-selective medium for the isolation and quantification of *Pseudomonas solanacearum*. *Bacterial Wilt Newsletter*, (10), 3-5.
- Fahy, P. C., & Persley, G. J. (1983). *Plant bacterial diseases: a diagnostic guide*. Academic Press, New York. 393pp.
- He, D. C., He, M. H., Amalin, D. M., Liu, W., Alvindia, D. G., & Zhan, J. (2021). Biological control of plant diseases: An evolutionary and eco-economic consideration. *Pathogens*, 10(10), 1311.
- Janse, J. D. (1988). A detection method for *Pseudomonas solanacearum* in symptomless potato tubers and some data on its sensitivity and specificity. *Eppo Bulletin*, 18(3), 343-351.
- Janse, J. D. (1996). Potato brown rot in Western Europe—history, present occurrence and some remarks on possible origin, epidemiology and control strategies. *Eppo Bulletin*, 26(3 4), 679-695.
- Kempe, J. (1983). Biological control of bacterial wilt of potatoes: attempts to induce resistance by treating tubers with bacteria. *Plant disease*, 67, 499-503.
- Khairy, A. M., Tohamy, M. R., Zayed, M. A., & Ali, M. A. (2021). Detecting pathogenic bacterial wilt disease of potato using biochemical markers and evaluate resistant in some cultivars. *Saudi Journal of Biological Sciences*, 28(9), 5193-5203.
- King, E. O., Ward, M. K., & Raney, D. E. (1954). Two simple media for the demonstration of pyocyanin and fluorescein. *The Journal of laboratory and clinical medicine*, 44(2), 301-307.
- Labuschagne, N., Pretorius, T., & Idris, A. H. (2011). Plant growth promoting rhizobacteria as biocontrol agents against soil-borne plant diseases. *Plant growth and health promoting bacteria*, 211-230.
- Lelliott, R. A., & Stead, D. E. (1987). *Methods for the diagnosis of bacterial diseases of plants*. Blackwell Scientific Publications. Vol. 2, Ed. T. F. Preece. Blackwell Scientific Publication 216 pp.
- Locci, R. (1989). *Streptomyces and related genera*. *Bergey's manual of systematic bacteriology*, 4, 2451-2508.
- Lwin, M. Y. I. N. T., & Ranamukhaarachchi, S. L. (2006). Development of biological control of *Ralstonia solanacearum* through antagonistic microbial populations. *International Journal of agriculture and biology*, 8(5), 657-660.
- Madigan, M.T.; Martiko, J.M. and Parker, J. (1997): *Antibiotics: Isolation and Characterization*. In Book *Biology of Microorganisms*. 8th edn. Prentice Hall International Inc. New Jersey, pp. 440- 442.
- Maji, S., & Chakrabartty, P. K. (2014). Biocontrol of bacterial wilt of tomato caused by '*Ralstonia solanacearum*' by isolates of plant growth promoting Rhizobacteria. *Australian Journal of Crop Science*, 8(2), 208-214.
- Mansfield, J., Genin, S., Magori, S., Citovsky, V., Sriariyanum, M., Ronald, P., ... & Foster, G. D. (2012). Top 10 plant pathogenic bacteria in molecular plant pathology. *Molecular plant pathology*, 13(6), 614-629.
- McCarter, S. M. (1976). Persistence of *Pseudomonas solanacearum* in artificially infested soils. *Phytopathology* 66: 998 – 1000.
- Michel, V. V., & Mew, T. W. (1998). Effect of a soil amendment on the survival of *Ralstonia solanacearum* in different soils. *Phytopathology*, 88(4), 300-305.
- Nion, Y. A., & Toyota, K. (2015). Recent trends in control methods for bacterial wilt diseases caused by *Ralstonia solanacearum*. *Microbes and environments*, 30(1), 1-11.
- Pérez-García, A., Romero, D., & De Vicente, A. (2011). Plant protection and growth stimulation by microorganisms: biotechnological applications of Bacilli in agriculture. *Current opinion in biotechnology*, 22(2), 187-193.
- Ran, L. X., Li, Z. N., Wu, G. J., Van Loon, L. C., & Bakker, P. A. H. M. (2005). Induction of systemic resistance against bacterial wilt in *Eucalyptus urophylla* by fluorescent *Pseudomonas* spp. *European Journal of Plant Pathology*, 113, 59-70.
- Rosyidah, A., Wardiyati, T., Abadi, A. L., & Maghfoer, M. D. (2013). Enhancement in effectiveness of antagonistic microbes by means of microbial combination to control *Ralstonia solanacearum* on potato planted in middle latitude. *J. Agrivita*, 35;174-183.
- Schaad, N. W., Jones, J. B., & Chun, W. (2001). *Laboratory guide for the identification of plant pathogenic bacteria* (No. Ed. 3). American Phytopathological society (APS press).
- Shaikh, S. S., Patel, P. R., Patel, S. S., Nikam, S. D., Rane, T. U., & Sayyed, R. Z. (2014). Production of biocontrol traits by banana field fluorescent *Pseudomonads* and comparison with chemical fungicide. *Indian J.Exp.Biol.*,52;917-92.

- Shaikh, S. S., Wani, S. J., & Sayyed, R. Z. (2018). Impact of interactions between rhizosphere and rhizobacteria: a review. *J Bacteriol Mycol*, 5(1), 1058.
- Shirling, E. T., & Gottlieb, D. (1966). Methods for characterization of *Streptomyces* species. *International journal of systematic bacteriology*, 16(3), 313-340.
- Soesanto, L., Manan, A., Wachjadi, M., & Mugiastuti, E. (2013). Ability test of several antagonists to control potato bacterial wilt in the field. *AGRIVITA, Journal of Agricultural Science*, 35(1), 30-35.
- Sunaina, V., Kishore, V., Shekhawat, G. S., & Kumar, M. (1997). Control of bacterial wilt of potatoes in naturally infested soils by bacterial antagonists / Bekämpfung der Bakterienwelke an Kartoffeln durch bakterielle Antagonisten in natürlich verseuchten Böden. *Zeitschrift Für Pflanzenkrankheiten Und Pflanzenschutz / Journal of Plant Diseases and Protection*, 104(4), 362–369. <http://www.jstor.org/stable/43215170>.
- Tan, S., Gu, Y., Yang, C., Dong, Y., Mei, X., Shen, Q., & Xu, Y. (2016). *Bacillus amyloliquefaciens* T-5 may prevent *Ralstonia solanacearum* infection through competitive exclusion. *Biology and Fertility of Soils*, 52, 341-351.
- Wagih, E. (1991). Neither Indole Acetic Acid nor Bacteriocin is Apparently Involved in the in vitro Antagonism between the Virulent and the Avirulent Strains of *Pseudomonas solanacearum*. *Journal of Phytopathology*, 132: 153-160. <https://doi.org/10.1111/j.1439-0434.1991.tb00106>.
- Wenneker, M., Verdel, M. S. W., Groeneveld, R. M. W., Kempenaar, C., Van Beuningen, A. R., & Janse, J. D. (1999). *Ralstonia (Pseudomonas) solanacearum* race 3 (biovar 2) in surface water and natural weed hosts: First report on stinging nettle (*Urtica dioica*). *European Journal of Plant Pathology*, 105, 307-315.
- Williams, S.T.; Goodfellow, M.; Andersons G. (1989): Genus *Streptomyces*. Waksman and Henrici 1943. 339AL. In *Bergey's Manual of Systematic Bacteriology*. Williams and Wilkins Company. Baltimore. Vol: 4. pp. 2452- 2492.
- Zhu, L., Wang, Y., Lu, J., Liu, S., Min, Y., Liu, X., ... & Zhou, R. (2021). Complete genome sequence of *Bacillus badius* NBPM-293, a plant growth-promoting strain isolated Wilkins Company. Baltimore. Vol: 4. pp. 2452- 2492.